

Donna Kubik Fall, 2006

- Scales of the Universe
- Patterns of Stars
- Earthly cycles
- Eclipses

- Scales of the Universe
- Patterns of Stars
- Earthly cycles
- Eclipses

## People-sized Spaces

- You
- Lecture Hall
- Faraday West
- NIU

- $5'9" = 1.75 \text{ m} = 1.75 \text{ x} 10^{\circ} \text{ m}$
- $14 \text{ m} = 1.4 \times 10^1 \text{ m}$
- $80 \text{ m} = 0.8 \times 10^2 \text{ m}$
- $2000 \text{ m} = 2.0 \times 10^3 \text{ m}$

## Map-Sized Spaces


- DeKalb (city)
- DeKalb County
- Illinois
- United States

- $5000 \text{ m} = 5 \text{ km} = 5 \text{ x} 10^3 \text{ m}$
- $30,000 \text{ m} = 30 \text{ km} = 3 \times 10^4 \text{ m}$
- $400 \text{ km} = 4 \times 10^5 \text{ m}$
- $5000 \text{ km} = 5 \times 10^6 \text{ m}$

## Space-Sized Spaces (Part 1)

- Earth
- Earth and Moon
- Earth to the Sun


- $13,000 \text{ km} = 1.3 \times 10^7 \text{ m}$
- $384,000 \text{ km} = 4 \times 10^8 \text{ m}$
- 150,000,000 km =  $1.5 \times 10^{11}$  m This is 1 Astronomical Unit (1 AU)



## Space-Sized Spaces (Part 2)

- Earth to the Sun
- Solar System

- 1 AU =  $1.5 \times 10^{11} \text{ m}$ 8 light-minutes
- 500 AU =  $0.8 \times 10^{14} \text{ m}$ 3 light-days




## Space-Sized Spaces (Part 3)

- Nearest Star
- Milky Way

- $300,000 \text{ AU} = 4 \times 10^{16} \text{ m}$ 
  - 4.3 light years = 4.3 ly
- $100,000 \text{ ly} = 10^{21} \text{ m}$





## Units often used in astronomy

- Astronomical Unit (AU)
- Light year (ly)
- Parsec (pc)

### Astronomical Unit


- Used to discuss distances across the Solar System
- 1 astronomical unit (AU) = average distance from the Earth to the Sun
- 1 AU = 150,000,000 km = 1.5x10<sup>8</sup> km



## Light Year


- Used to discuss distances to the stars
- 1 light year (ly) = distance that light travels in one year
- 1 ly =  $9.46 \times 10^{12}$  km
- 1 ly = 63,000 AU
- Note: Remember light travels at ~3x10<sup>8</sup> m/s

Distance to the closest star (Proxima Centauri) = 4.2 ly



### Parsec

- Another unit used to discuss distances to the stars
- 1 parsec = 1 pc = 3.26 ly




- Scales of the Universe
- Patterns of Stars
- Earthly cycles
- Eclipses

## Constellations





- Constellations are patterns formed by bright stars
- The celestial sphere is divided into 88 unequal regions.
- These regions are what astronomers call constellations
- Astronomers use the term to describe an entire region of the sky and all the objects in that region

• We see the projected pattern of stars of varying distances




The stars in a constellation are not even near each other!





## Map of the constellations




- Historically, constellations
   were groupings of stars that
   were thought to outline the
   shape of something, usually
   with mythological
   significance.
- There are 88 recognized constellations, with their names tracing as far back as Mesopotamia, 5000 years ago.
- In modern astronomy, the significance of constellations is no longer mythological, but practical: constellations define imaginary regions of the sky
- Just as the individual states each define an imaginary region of the United States.

- Thus, to say that an object (planet, Moon, cluster, nebula, etc) is "in" the constellation Scorpius is to partially locate the planet on the celestial sphere, just as saying that Knoxville is in Tennessee is to partially locate the city on the surface of the Earth.
- Like states, modern constellations have irregular boundaries that have been agreed upon for various reasons, perhaps not always completely logical.

## Using constellations as landmarks

- So you could say:
- "At dawn on Jan 25, the Moon will be in Scorpius."
- "Jupiter will be in Libra."




### Star names

- Stars are generally designated by a Greek letter and the name of the constellation in which they reside.
- Usually  $\alpha$  is the brightest star in the constellation,  $\beta$  is the second brightest, etc
- Sirius is the brightest star in the night sky



### Constellations

 Stars move on geologic timescales, so a constellation's pattern will change as is shown on the right for the Big Dipper




2000 A.D

 The patterns are unchanging in ~10,000 year timescales




50,000 A.D

 BUT, their position in sky varies with season (as shown in next slide)




100,000 A.D

## Different seasons, different constellations




### Summer constellations

Cygnus (The Swan)






## Summer triangle




### Winter Constellations



#### Orion



## Winter triangle




### Constellations

 "The constellations have such intriguing names - somehow we expect the books to show us groups of stars in the shape of a Lion, a Whale, Twins, etc. But they show us nothing of the sort. This book serves to remedy the situation."


- ~H A Rey


## H. A. Rey

- H. A. Rey "The Stars, A New Way to See Them"
- H. A. Rey also wrote the "Curious George" books!




## Traditional vs. H. A. Rey





## Map of the constellations



http://www.astro.wisc.edu/~dolan/constellations/constellationjavalist.html

### Constellations

- Planetarium programs often give the option to display constellations using Rey's or astronomical stick figures.
- The free program on Sky and Telescope's website uses Rey's figures! <a href="http://skyandtelescope.com/">http://skyandtelescope.com/</a>




### Polaris - The North Star

Polaris is almost directly overhead at the North Pole.



- Polaris is at the north horizon at the Equator.
- Polaris is not visible south of the Equator.
- Other stars appear to move due to the Earth's rotation.




### Polaris - The North Star

Light rays from Polaris


 Polaris is always located above the northern horizon at an angle equal to your latitude

 De Kalb's latitude is about 42° N




## Rising and Setting Stars

- Stars appear to "move" in circles about Polaris.
- This is because the Earth is rotating about the geographic pole.
- It takes 1 day to complete the circle.
- Stars that never rise or set are called *circumpolar* stars




## Star trails in the northern sky

- As the earth spins on its axis, the sky seems to rotate around us.
- This motion produces the concentric trails traced by the stars in this time exposure of the night sky.
- The north celestial pole (NCP) is at the center.



## Star trails in the northern sky

- The very short bright trail near the NCP was made by Polaris, commonly known as the North Star.
- So this is proof that Polaris is not *exactly* due north.



## Star trails in the northern sky


 More proof that Polaris is not exactly due north.



Star trails in the southern sky

 While the bright star Polaris lies conveniently close to the North Celestial Pole, no bright star similarly marks the pole in the South.


 Still, the South Celestial Pole is easily identified in the picture as the point in the sky at the center of all the star trail arcs.



# Calculate time of exposure from star trails

- Measure the angle subtended by the star trails (N deg)
- Then the time of the exposure was:

(N deg)(1 day/360 deg)(24 hours/day) = X hours



# Calculate time of exposure from star trails

 The angle subtended by each star trail is ~20 degrees, so the time of the exposure was:

(20 deg)(1 day/360 deg)(24 hours/day) = 1.3 hours

### Star trails at mid-latitudes

- Star trails from a lower latitude
- Notice that the trails set beneath the horizon
- These trails were made by stars that are NOT circumpolar stars




## Star trails at mid-latitudes

 Star trails from a lower latitude (trails set beneath the horizon)



# Compare rising and setting at high and low latitudes





# Equatorial mount on Celestron14 as in Davis Hall Observatory, NIU


#### To Polaris






## Celestial Sphere

- The stars can be mapped onto a sphere.
- This sphere is called the celestial sphere
- Positions correspond to points over locations on earth.




## Celestial Sphere

- The North and South
   Celestial Poles correspond
   to the Geographic North and
   South Poles of Earth
- The Celestial Equator corresponds to the Earth's Equator



## Celestial Sphere

 The coordinates of objects on the celestial sphere are analogous to longitude and latitude



## Longitude and latitude

DeKalb, IL

Latitude: 41.93 N

Longitude: 88.75 W




### Latitude and declination

#### Latitude

 Latitude measures the number of degrees north or south of the Equator. DeKalb is at 41.93° N latitude.

#### **Declination**


 Stars have a north-south position called the declination which is similar to the idea of latitude



### Latitude and declination

#### **Declination**

- The declination of a star is the number degrees north or south of the Celestial Equator
- Polaris has a declination of 90° N.
- Stars over the equator have a declination of 0°.




## Longitude and Right Ascension

#### Longitude

 Longitude measures the number of degrees east or west of the Meridian. DeKalb is at 88.75° W longitude.

#### Right Ascension


 Stars have an east-west position called the *right* ascension which is similar to the idea of longitude.



## Longitude and Right Ascension

#### Right Ascension


- The Vernal (spring) equinox is used to define zero hours right ascension.
- Right ascension is the number of hours, minutes, seconds east or west of the Vernal equinox



## Longitude and Right Ascension


#### Right Ascension

 Vernal equinox is the point of intersection of the celestial equator and the ecliptic, where the sun crosses the celestial equator moving northward.




### Star and Planet locator


- A Star and Planet Locator is a map of the celestial sphere
- Other names for a Star and Planet Locator are
  - Planisphere
  - Star Finder




### Star and Planet Locator

- The earth turns under the stars, so the stars turn on the Star and Planet Locator.
- Positions correspond to points over locations on earth (to positions on the celestial sphere)






### Star and Planet Locator



### Horizon


- The earth blocks stars too far to the south.
- The *horizon* is the line of the ground for an observer.
- A star finder provides a

   cover that act as the horizon.
  - You use a different cover depending on your latitude




## Line up the time to the date


 The wheel turns to set the day and time for the observer.




 The Star and Planet Finder mimics the Earth's daily rotation and yearly revolution about the Sun




- Stars "move" east to west over the course of one night in circle about the North Celestial Pole
- As you spin the disk CCW, see how new stars 'appear' from under the eastern horizon and stars 'disappear' under the western horizon
- This illustrates the rising and setting of stars




- Stars "move" east to west by 2 hours per month and "return" to the same position after one year
- This is due to the difference between the solar and sidereal day (described a few slides later)




- Notice that the days are separated by 1 degree.
- So for the same time on successive nights, the stars' positions have shifted
- Stars rise ~4 minutes earlier each day
- (1 day)(24 hours/day)(60 min/hour)=4 min




 The ecliptic is the plane described by the Earth's path around the Sun




- The ecliptic and celestial equator are different circles tilted 23.5 degrees with respect to each other on the celestial sphere
- This is because the earth's axis of rotation is tilted 23.5 degrees from the plane of the ecliptic




 The ecliptic is indicated by dashes on a Planisphere




 Each planet, including the Earth's, axis of rotation is inclined to the plane of the ecliptic.



 The orbit of the Moon and the orbits of many of the planets lie near the plane of the ecliptic



Inner planets



Inner and outer planets

 The Plane of the Ecliptic is illustrated in this Clementine star tracker camera image which reveals (from right to left) the Moon lit by Earthshine, the Sun's corona rising over the Moon's dark limb, and the planets Saturn, Mars, and Mercury.



## Stars vs. Planets

 How can you tell the difference between a star and a planet?



- How can you tell the difference between a star and a planet?
- The differences seen in this image are enumerated on the following slides



- Image of Jupiter and Saturn
- Among the background stars are the familiar Pleiades (above right) and V-shaped Hyades (below left) star clusters
- Animated sequence combining 23 pictures taken at approximately 2 week intervals from June 2000 through May 2001.



#### Stars

Stars are very numerous.

#### **Planets**

• Planets are few in number (5 are visible to the unaided eye).



#### Stars

• Stars are "fixed" relative to each other.

#### **Planets**

 Planets "wander" relative to the fixed stars. So they are not in the same location each night nor in the same position year to year



#### Stars

 Stars have a wide range of declination and right ascension

#### **Planets**

 Must be on (or very near) the ecliptic



#### Stars

#### They produce their own light independent of the Sun's location.

Stars twinkle

They are very far away

#### **Planets**

- Their brightness <u>does</u> depend on the Sun's location.
  - Planets don't twinkle


• Relative to the stars, they are near to Earth.

# Chapter 1 Discovering the Night Sky

- Scales of the Universe
- Patterns of Stars
- Earthly cycles
- Eclipses


## Seasons

- The seasons are caused by the tilt of the Earth's axis of rotation
- The seasons are NOT due to the nearness of the Earth to the Sun
- In fact, the Earth is closest to the Sun around January 3 of each year!



## Seasons

- In summer, the Sun is more directly overhead in the northern hemisphere.
- The Sun is also hitting us for more hours each day.
- Hence it's warmer in the summer.
- Seasons <u>Movie</u>



http://sealevel.jpl.nasa.gov/overview/climate-spherical.html

#### Diurnal and Annual Motion


#### Diurnal (daily) motion

- Earth spins once per day.
- Such motion is called rotation
- Causes stars to appear to move east to west over the course of one night in a circle about the North Celestial Pole.

#### Annual (yearly) motion

- Earth orbits the Sun once per year
- Such motion is called revolution
- Causes each star to rise approximately 4 minutes earlier each night than it did the night before






Sidereal refers to stars

Solar refers to the Sun





Because the Earth moves along its orbit around the Sun, the Earth must rotate an extra degree (~361 deg) to get from one local noon to the next.

This extra degree of rotation corresponds to four minutes of time.


Sidereal day = 23h 56m 4.091s

Solar day = 24 hours

# Synodic vs. sidereal month

- A *synodic* month (29.5 days) is the time it takes the Moon to complete a full cycle of phases.
- A *sidereal* month (27.3 days) is the time it takes for the Moon to complete one 360 degree orbit about the earth.
- The difference is due to the earth's constant motion in its orbit about the sun; the Moon must travel through >360 degrees to get from one new Moon to the next.

# \* Sidereal vs. synodic month




The sidereal month is the time the Moon takes to complete one full revolution around the Earth with respect to the background stars.

However, since the Earth is constantly moving along its orbit around the Sun, the Moon must travel slightly more than 360 degrees to get from one new moon to the next.

Thus the *synodic* month is longer than the sidereal month.



# → The moon rises ~50 min later each day




The Moon takes  $\sim 29.5$  days to orbit the earth.

This corresponds to  $\frac{360^{\circ}}{29.5 days} = \frac{12.2^{\circ}}{day}$ .  $\rightarrow \left(\frac{12.2^{\circ}}{day}\right)\left(\frac{24hours}{360^{\circ}}\right)\left(\frac{60\min}{hour}\right) = 48.8\min$ 

So the Moon rises  $\sim 48.8$  minutes later each day.



→ The moon rises ~50 min later each day



The Moon takes  $\sim 29.5$  days to orbit the earth.

This corresponds to  $\frac{360^{\circ}}{29.5 \, days} = \frac{12.2^{\circ}}{day}$ .  $\rightarrow \left(\frac{12.2^{\circ}}{day}\right)\left(\frac{24hours}{360^{\circ}}\right)\left(\frac{60 \text{ min}}{hour}\right) = 48.8 \text{ min}$ 

So the Moon rises  $\sim 48.8$  minutes later each day.


#### Moon & month

- The approximately 4 weeks it takes the Moon to complete one orbit inspired the concept of our month
- Month lengths formerly followed motions of the moon. But our calendar's irregular distribution of days in the various months no longer corresponds to either the lunar cycle or to that of any other object in the heavens.



## Time of moonrise


• The moon rises about 50 minutes later each day than the day before.



• Like the Earth, the Moon is a sphere which is always half illuminated by the Sun, but as the Moon orbits the Earth we get to see more or less of the illuminated half.



 This image was taken by NEAR as it flew by the Earth-Moon system in January 1998.




• As the Moon orbits the Earth we get to see more or less of the illuminated half.



Waxing crescent

Waxing gibbous


Third quarter




First quarter

Waning gibbous


Waning crescent





# The "age" of the moon

- The time in days counted from the time of new Moon is called the moon's age
- The age of the moon is measured in *days*.



#### Blue Moon

- Because the cycle of the phases is shorter than most calendar months, the phase of the Moon at the very beginning of the month usually repeats at the very end of the month.
- When there are two full Moons in a month (which occurs, on average, every 2.7 years), the second one is called a "Blue Moon".

#### Lunation

 Our Moon's appearance changes nightly. This timelapse sequence shows what the Moon looks like during a lunation, a complete lunar cycle.



# Things that change during a lunation

- The apparent size changes
- The phase changes
- We can see a different area (not always the same 50%)



# Apparent size of the Moon changes

 Due to the elliptical shape of the Moon's orbit, the apparent size of the Moon's disk changes as its distance from Earth varies



# Apparent size of the Moon changes

 Note that the closest and farthest points do not always occur at the same phases



# 59% of the Moon's surface is observable from Earth


- Although the Moon's rotation on its axis is synchronously locked with its revolution around Earth, *librations* permit a terrestrial observer to see slightly differing halves of the Moon's surface at different times.
- This means that a total of, not 50%, but 59% of the Moon's surface can be observed from Earth.




#### Libration

- A libration (from the Latin verb libro -are "to balance, to sway", cf. libra "scales") is a very slow oscillation, real or apparent, of a satellite as viewed from the larger celestial body around which it revolves.
- The term usually refers to the apparent movements of the Moon relative to Earth, which can be compared to the rocking of a pair of scales about the point of balance.
- Note same origin as that of the constellation Libra

## The Constellation Libra





# 3 types of libration

- Libration in latitude is a consequence of the Moon's axis of rotation being slightly inclined to the normal to the plane of its orbit around Earth. Its origin is analogous to the way in which the seasons arise from Earth's revolution about the Sun.
- Libration in longitude is a consequence of the Moon's orbit around Earth being somewhat eccentric, so that the Moon's rotation sometimes leads and sometimes lags its orbital position.
- Diurnal libration. This is a consequence of Earth's rotation. This carries an observer first to one side and then to the other of the straight line joining Earth's center to the Moon's center, allowing the observer to look first around one side of the Moon and then around the other.


## Planets have phases, too

 Voyager 2 spacecraft camera captured Neptune and Triton together in crescent phase in 1989.



## Planets have phases, too


 The inferior planets exhibit a complete set of phases (just like the Moon) as viewed from the Earth



Aspects & Phases of the Inferior Planets

### Planets have phases, too

- The superior planets do not exhibit a full range of phases
- They are always gibbous or full




Aspects & Phases of the Superior Planets

# Chapter 1 Discovering the Night Sky

- Scales of the Universe
- Patterns of Stars
- Earthly cycles
- Eclipses

### **Eclipses**



#### How Eclipses Work

- The Moon or Earth can block the sun.
- But eclipses cannot happen every full or new Moon

- Lunar eclipse can occur Full Moon

  New Moon
  Solar eclipse can occur
- The Moon's orbit is not in the same plane as the Sun; it is tilted 5 degrees out of the ecliptic
- Eclipse season is twice a year.



Eclipse season is twice a year Full Lunar eclipse can occur 5° Full Line of nodes 5° New Solar eclipse can occur No eclipse possible Sun 5° Earth's Solar eclipse can occur New orbit 5° No eclipse possible Full Lunar eclipse can occur Full

### Kinds of Eclipses

Solar Eclipse

• Lunar Eclipse

Moon's shadow on the earth

Earth's shadow on the moon.

Only occurs at a new moon.

Only occurs at a full moon.

### Kinds of Eclipses

Solar Eclipse

Partial - crescent sun remains

Total - all of the sun is blocked

Annular - a ring of the sun remains

• Lunar Eclipse

Partial - moon is partially dark

Total - moon is dark red or copper-colored (explained in a few slides)

Earth's atmosphere passes some light.

### Kinds of Eclipses

Solar Eclipse

• Lunar Eclipse

Maximum duration of totality:


7.5 minutes

Maximum duration:

1 hour 47 minutes


## Why is the sky blue and the sunset red?

- A clear, cloudless day-time sky is blue, because molecules in the air scatter blue light from the sun more than they scatter red light.
- This scattering is called Rayleigh scattering and is proportional to  $1/\lambda^4$
- So it is a 10 times larger effect for blue (300 nm) than for red (700 nm) light




## Why is the sky blue and the sunset red?

- When we look towards the sun at sunset, the light must travel through much more atmosphere than when the sun is more directly overhead
- So we see red and orange colors, because more of the blue light has been scattered out and away from the line of sight



# Why does the Moon appear reddish during lunar eclipse?

- During a lunar eclipse, this red light is refracted by the Earth's atmosphere and can reflect off the Moon.
- This makes the Moon look reddish



## Why does the Moon appear reddish during lunar eclipse?


- The red tint of the eclipsed Moon is created by sunlight first passing through the Earth's atmosphere, which preferentially scatters blue light (making the sky blue) but passes and refracts red light, before reflecting back off the Moon.
- Differing amounts of clouds and volcanic dust in the Earth's atmosphere make each lunar eclipse appear differently.



# Why does the Moon appear reddish during lunar eclipse?

 Before and after totality, can see some sunlight reflected off the moon

 During totality, no direct sunlight hits the moon, only the red light refracted through the earth's atmosphere



### Viewing Eclipses

- Never look directly at the sun, even at a partial eclipse
- View sun through #14 welder's glass or special mylar filters.
- It's ok to look at lunar eclipses





#### Eclipse Prediction

- The cycles of the sun and moon cause eclipses in a predictable pattern.
- The same eclipse occurs every 18 years the saros cycle.
- Last total solar eclipse was December 4, 2002 in Africa, Indian Ocean and Australia.
- Next partial solar eclipse in DeKalb is March 19, 2007.
- Next total solar eclipse in the US is August 21, 2017.

### Science during eclipses

- Eclipses are fun
- Eclipses also provide a golden opportunity to learn more about the sun.
- The hot gases surrounding the Sun can be observed, providing information about its
  - temperature
  - chemistry
  - atmospheric activity.



### Total Eclipse of the Sun







Total Eclipse of the Heart

Bonnie Tyler

http://www.eso.org/outreach/info-events/eclipse99/report-hq.html
Link to video of total eclipse, August 11, 1999, Bavaria, Germany
The duration of totality 2 min 17 sec.

